

How Old is the Universe?

J. W. Moffat

Department of Physics

University of Toronto

Toronto, Ontario M5S 1A7, Canada

ABSTRACT

A dynamical model of the decaying vacuum energy is presented, which is based on Jordan-Brans-Dicke theory with a scalar field ϕ . The solution of an evolutionary differential equation for the scalar field ϕ drives the vacuum energy towards a cosmological constant at the present epoch that can give for the age of the universe, $t_0 \sim 13.5$ Gyr for $\Omega_0 = 1$, which is consistent with the age of globular clusters.

UTPT-94-34

e-mail: moffat@medb.physics.utoronto.ca

1. Introduction

The age of the universe problem is a well-known quandary of the spatially homogeneous models. Estimates of the age of the Galaxy based on radioactive dating, the age of globular clusters and other methods give the age of the universe within 10 to 20 Gyr¹. At the same time, in order for a matter dominated FRW model with the density parameter $\Omega_0 = 1$ to yield $t_0 \geq 10$ Gyr requires $h \leq 0.65$, where h is related to the observationally established value of the Hubble constant through $H_0 = 100h\text{kms}^{-1}\text{Mpc}^{-1}$. A value of $h \geq 0.65$ produces the age of the universe problem. Recent measurements using the Canada-France-Hawaii Telescope² and the Hubble Space Telescope³ have given $H_0 = 87 \pm 7\text{kms}^{-1}\text{Mpc}^{-1}$ and $80 \pm 17\text{kms}^{-1}\text{Mpc}^{-1}$, respectively. These results imply an age for the universe of 7×10^9 yrs and 8×10^9 yrs, respectively, in the standard big bang model with $t_0 = 2/3 \times 1/H_0$. These results have produced a new age of the universe crisis.

The inflationary paradigm⁴ has motivated the need to have $\Omega_0 = 1$. Also, dynamical estimates of the FRW density parameter Ω_0 give results for observations over large scales (> 20 Mpc, say ~ 100 Mpc) indicating the existence of a less clustered component with a contribution perhaps as high as $\Omega_{\sim 100} \simeq 0.8 \pm 0.2$ ⁵.

The standard solution to the age of the universe problem is based on the postulate that the cosmological constant Λ is non-zero⁶. For $\Omega_{\text{matter}} = 0.2$ and $\Omega_{\text{vacuum}} = 0.8$, corresponding to $\Omega = \Omega_{\text{matter}} + \Omega_{\text{vacuum}} = 1$, the expansion age is $t_0 = 13.5$ Gyr, which is consistent with the other estimates of the age. But the cosmological constant is very small, $\Lambda < 10^{-46} \text{GeV}^4$, so its use to obtain a consistent age of the universe introduces a severe fine-tuning problem.

The cosmological constant provides a source of negative pressure, producing an acceleration of the expansion of the universe that negates the deceleration of the expansion

caused by matter and radiation. This makes the universe appear younger than it is to an observer who assumes that the cosmological constant is zero. If $H_0 t_0$ exceeds unity, it means the expansion has been accelerating rather than decreasing, and within the standard model this would imply that the active gravitational mass density is negative, which would happen if there were an appreciable positive cosmological constant.

In Einstein's field equations:

$$G_{\mu\nu} + \Lambda_B g_{\mu\nu} = 8\pi G T_{\mu\nu}, \quad (1.1)$$

Λ_B is called the “bare” cosmological constant, while a second cosmological constant enters through the vacuum energy density:

$$T_{V\mu\nu} = -\rho_V g_{\mu\nu} = -\frac{\Lambda_V}{8\pi G} g_{\mu\nu}. \quad (1.2)$$

These two constants can be combined through an effective cosmological constant:

$$\Lambda = \Lambda_B + \Lambda_V. \quad (1.3)$$

Why Λ has the incredibly small value, $\Lambda < 10^{-46} \text{GeV}^4$, today is known as the cosmological constant problem⁷.

We shall seek an explanation for the smallness of Λ at the present epoch, and the fact that it has a critical value needed to resolve the age of the universe problem by using the Jordan-Brans-Dicke (JBD) action^{8–12}. In JBD, the kinetic energy can produce a negative pressure, a fact emphasized recently by Levin and Freese¹³ in connection with inflationary theory. The nonminimal coupling of the scalar field to gravity allows for a negative pressure associated with the kinetic energy of the field. By assuming that Λ is spacetime dependent and equating it to the kinetic energy in the JBD theory, we find that the evolutionary equation for the scalar field in an FRW universe, produces an attractor mechanism that

drives Λ towards a minimum of the potential. This damping mechanism can give a small constant value for Λ today that can resolve the age of the universe problem.

We shall assume that $\Lambda_B = 0$ and that the vacuum energy is dominated by the scalar field ϕ . Unbroken supersymmetry can “protect” the constant Λ_B from becoming non-zero⁷, but in the real universe supersymmetry will be broken at an energy > 1 TeV. We do not know of any other symmetry that guarantees that Λ_B remains zero.

2. Tensor-Scalar Theory

In the Jordan-Fierz conformal frame the Jordan-Brans-Dicke action for a massless tensor-scalar theory takes the form:

$$\tilde{S} = \frac{1}{16\pi} \int d^4x \sqrt{-\tilde{g}} \left[\psi \tilde{R} - \frac{\omega(\psi)}{\psi} \tilde{g}^{\mu\nu} \psi_{,\mu} \psi_{,\nu} \right] + S_m, \quad (2.1)$$

where $\tilde{R} = \tilde{g}^{\mu\nu} \tilde{R}_{\mu\nu}$ denotes the curvature scalar in terms of the physical metric tensor $\tilde{g}_{\mu\nu}$, $\omega(\psi)$ denotes the JBD ψ -dependent coupling parameter and S_m denotes the action for matter. In the “Einstein frame” the action reads:

$$S = \frac{1}{16\pi G} \int d^4x \sqrt{-g} (R - 2g^{\mu\nu} \phi_{,\mu} \phi_{,\nu}) + S_m, \quad (2.2)$$

where $g_{\mu\nu}$ denotes the “Einstein” metric tensor conformally related to the physical Jordan-Fierz one by the transformation:

$$\tilde{g}_{\mu\nu} = C^2(\phi) g_{\mu\nu} \quad (2.3)$$

and $R = g^{\mu\nu} R_{\mu\nu}$.

Let us now define the scalar field:

$$\Lambda(x) = g^{\mu\nu} \phi_{,\mu} \phi_{,\nu}, \quad (2.4)$$

and identify $\Lambda(x)$ as a variable cosmological “constant”. Then, Eq.(2.2) becomes:

$$S = \frac{1}{16\pi G} \int d^4x \sqrt{-g} (R - 2\Lambda) + S_m. \quad (2.5)$$

The field equations in the Einstein conformal frame are:

$$R_{\mu\nu} = 2\Lambda_{\mu\nu} + 8\pi G(T_{\mu\nu} - \frac{1}{2}g_{\mu\nu}T), \quad (2.6)$$

$$\square_g \phi = -4\pi G a V(\phi) T, \quad (2.7)$$

where $T_{\mu\nu}$ denotes the Einstein stress-energy tensor, $\square_g = g^{\mu\nu} \nabla_\mu \nabla_\nu$ and ∇_μ is the covariant derivative with respect to the $g_{\mu\nu}$ connection. Moreover,

$$\Lambda_{\mu\nu} = \phi_{,\mu} \phi_{,\nu} \quad (2.8)$$

and

$$a(\phi) \equiv \frac{\partial V(\phi)}{\partial \phi} \quad (2.9)$$

is the gradient of the coupling function $V(\phi)$ where

$$V(\phi) = \ln C(\phi). \quad (2.10)$$

3. Tensor-Scalar Cosmology with Varying Λ

The metric takes its usual FRW form:

$$ds^2 = dt^2 - R^2(t) \left[\frac{dr^2}{1 - kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right], \quad (3.1)$$

where $k = +1, 0, -1$ denotes the sign of the spatial curvature. The matter distribution is described by a perfect fluid form:

$$T^{\mu\nu} = (\rho + p)u^\mu u^\nu - pg^{\mu\nu} \quad (3.2)$$

with $g_{\mu\nu}u^\mu u^\nu = \tilde{g}_{\mu\nu}\tilde{u}^\mu\tilde{u}^\nu = 1$, $\rho = C^4\tilde{\rho}$ and $p = C^4\tilde{p}$. The Jordan-Fierz variables $\tilde{t}$ (physical cosmic time) and $\tilde{R}$ are given by

$$d\tilde{t} = C(\phi(t))dt, \quad (3.3)$$

$$\tilde{R}(\tilde{t}) = C(\phi(t))R(t). \quad (3.4)$$

The Einstein equations (2.6) give

$$\frac{\ddot{R}}{R} = -\frac{4\pi}{3}G(\rho + 3p) - \frac{2\Lambda}{3}, \quad (3.5)$$

$$\left(\frac{\dot{R}}{R}\right)^2 + \frac{k}{R^2} = \frac{8\pi G\rho}{3} + \frac{\Lambda}{3}, \quad (3.6)$$

where $\Lambda = (\dot{\phi})^2$ and $\dot{\phi} = d\phi/dt$. The scalar field equation is

$$\ddot{\phi} + (\dot{\phi})^2 + 3\frac{\dot{R}}{R}\dot{\phi} = -4\pi Ga(\phi)(\rho - 3p). \quad (3.7)$$

The conservation equation reads:

$$d(\rho R^3) + p dR^3 = (\rho - 3p)R^3 dV(\phi). \quad (3.8)$$

We introduce the evolution parameter:

$$\tau = \ln R + \text{const}, \quad (3.9)$$

and $d\tau = H dt$ where $H = \dot{R}/R$ is the Hubble parameter in the Einstein frame. We also have

$$\dot{\phi} = H \frac{d\phi}{d\tau}. \quad (3.10)$$

4. Scalar Field Evolution Equation

The evolution equation for the scalar field ϕ for $k = 0$ takes the form:

$$\frac{2}{3 - \phi'^2}\phi'' + (1 - \theta)\phi' = -(1 - 3\theta)a(\phi), \quad (4.1)$$

where $\phi' = d\phi/d\tau$ and $\theta = \tilde{p}/\tilde{\rho}$ is considered to be either a constant or a known function of τ .

The trajectories in the $\phi, \dot{\phi}$ phase space associated with the solutions of Eq.(4.1) have been studied extensively by Belinskii et al.,¹⁴ and by other authors^{15–18}. They showed that the point of origin, $\phi, \dot{\phi} = (0, 0)$, is a point of stable focus and is the only point of equilibrium, while other singular points can lie only at infinity.

Eq.(4.1) describes the motion of a “ ϕ ” particle in a curved spacetime with a limiting speed equal to $\sqrt{3}$, a friction term proportional to the velocity, ϕ' , and a force term proportional to the gradient of the potential, $V(\phi)$. For a dominant energy condition, the physical energy density $\tilde{\rho}$ is positive and $-1 < \theta < +1$, so that the second term in (4.1) always damps the motion of the particle. For an inflationary era the force term on the right-hand side of (4.1) is negative ($\theta = -1$); it nearly vanishes for the radiation dominated era ($\theta \approx 1/3$) and is negative for the matter dominated one ($\theta \approx 0$).

In the radiation dominated era, Eq.(4.1) gives to first approximation:

$$\frac{2}{3 - \phi'^2} \phi'' + \frac{2}{3} \phi' = 0. \quad (4.2)$$

A solution of (4.2) is¹⁸

$$\phi(\tau) = \phi_\infty - \sqrt{3} \ln[B \exp(-\tau) + (1 + B^2 \exp(-2\tau))^{1/2}], \quad (4.3)$$

where

$$B = \frac{\phi'_0/\sqrt{3}}{\sqrt{1 - \phi_0'^2/3}}. \quad (4.4)$$

The total displacement of a particle between $\tau = 0$ and $\tau = \infty$ is given by

$$\Delta\phi = \phi_\infty - \phi_0 = \frac{1}{2} \sqrt{3} \ln \frac{1 + \phi'_0/\sqrt{3}}{1 - \phi'_0/\sqrt{3}}. \quad (4.5)$$

For non-relativistic motion, $\phi'_0 \ll \sqrt{3}$, we get

$$\Delta\phi \approx \phi'_0. \quad (4.6)$$

Thus, upon entering the radiation era the ϕ particle will fall down the potential $V(\phi)$ by a small amount.

For the matter dominated era with $\theta = \tilde{p}/\tilde{\rho} \ll 1$, the evolution equation becomes

$$m(\phi')\phi'' + \phi' = -a(\phi), \quad (4.7)$$

with the mass:

$$m(\phi') = \frac{2}{3 - \phi'^2}. \quad (4.8)$$

Here $m(\phi')$ has the non-relativistic limit $m(\phi') \approx 2/3$. The ϕ particle enters the matter era with an approximately zero initial velocity and $\sqrt{3}$ is a critical value for $|\phi'|$. The long-term behaviour of the particle depends dominantly on the shape of the potential $V(\phi)$ and *the particle ends up near a local minimum of $V(\phi)$* .

The behavior of the solution to Eq.(4.7) can be analyzed for the simple case of a parabolic potential¹⁸:

$$V(\phi) = \frac{1}{2}\kappa\phi^2, \quad (4.9)$$

where κ is the curvature:

$$\kappa(\phi) = \frac{\partial a(\phi)}{\partial \phi} = \frac{\partial^2 V}{\partial \phi^2} \quad (4.10)$$

calculated at the bottom of the well. For the potential (4.9), $\kappa(\phi) = \kappa$ is a constant. The solution to Eq.(4.7) has a critical value, $\kappa = 3/8$, and for $0 < \kappa < 3/8$ the motion is monotonic and is a linear combination of decreasing exponentials with positive coefficients.

For $\kappa > 3/8$, the motion is a damped oscillatory one and when $\kappa \rightarrow 3/8$, the solution tends towards the critically damped solution:

$$\phi(\tau) = \phi_R(1 + \frac{3}{4}\tau)\exp(-\frac{3}{4}\tau), \quad (4.11)$$

where ϕ_R is the value of ϕ at the end of the radiation era.

We expect that after a long time the particle is forced towards a minimum of $V(\phi)$ (near where $a(\phi)$ vanishes). Thus, an *attractor mechanism* forces the variable cosmological term Λ towards small or zero values.

5. Present Value of the Cosmological Constant

In the matter dominated era, we can rewrite Eq.(3.6) in terms of the Jordan-Fierz variables and $\phi(\tau)$ (recall that $\Lambda = (\dot{\phi})^2$):

$$\tilde{\Omega}_M = \frac{(1+a^2)(1-\phi'^2/3)}{(1+a\phi')^2} + \frac{(1+a^2)k}{\tilde{H}^2 \tilde{R}^2}, \quad (5.1)$$

where

$$\tilde{\Omega}_M = \frac{8\pi\tilde{G}\tilde{\rho}_M}{3\tilde{H}^2}. \quad (5.2)$$

Here, $\tilde{\rho}_M$ is the matter energy density and

$$\tilde{G}(\phi) = GC^2(\phi)[1+a^2(\phi)]. \quad (5.3)$$

The observational limits on a are (1σ level):

$$a_{\text{solar system}}^2 < 0.01. \quad (5.4)$$

Then (5.1) gives approximately for $k = 0$ at the present epoch:

$$\tilde{\Omega}_0 = \tilde{\Omega}_{0M} + \tilde{\Omega}_{0V} \approx 1, \quad (5.5)$$

where from (3.3), (3.4) and (3.10), we have

$$\tilde{\Omega}_{0V} = \frac{\phi_0'^2}{3} = \frac{(d\phi_0/d\tilde{t})^2}{3\tilde{H}_0^2}, \quad (5.6)$$

which can be written as

$$\tilde{\Omega}_{0V} = \frac{8\pi G\tilde{\rho}_{0V}}{3\tilde{H}_0^2} = \frac{\tilde{\Lambda}_0(\phi_0)}{3\tilde{H}_0^2}, \quad (5.7)$$

where $\tilde{\rho}_{0V}$ denotes the vacuum energy density at the present epoch.

We now have a scenario in which the scalar velocity is damped out to its present value ϕ_0' at a minimum of the potential $V(\phi)$. This corresponds to having an initially large cosmological term $\tilde{\Lambda}(x)$ during the inflationary period in the early universe, which is damped out in the matter dominated era to its present value $\tilde{\Lambda}_0$.

In the standard picture of inflation, the cosmic inflation is driven by the potential for the scalar field ϕ (i.e., $\phi'' \approx 0$ in Eq.(4.1)), whereas we have the opposite situation in which negative pressure is produced by the kinetic energy of the JBD theory. We shall assume that the standard potential-driven scenario will hold at near Planckian times, producing an inflationary expansion. Then, as the universe enters the non-inflationary era, the ϕ'' term in (4.1) becomes important and we have the scenario which drives Λ to small or zero values as ϕ' approaches the minimum of $V(\phi)$.

6. The Age of the Universe

For $k = 0$ and a dust-dominated universe, the present age of an object that formed at a redshift z_c is⁷:

$$t_c(z_c) = \frac{2}{3} \left(1 + \frac{\Omega_M}{\Omega_V}\right)^{1/2} H_0^{-1} \left\{ \sinh^{-1} \left(\frac{\Omega_V}{\Omega_M} \right)^{1/2} - \sinh^{-1} \left[\left(\frac{\Omega_V}{\Omega_M} \right)^{1/2} (1 + z_c)^{-3/2} \right] \right\}. \quad (6.1)$$

If we choose $z_c = 4$ and $\Omega_V/\Omega_M = 9$ ($\Omega_M = 0.1$), then this yields $t_0 = 1.1 \times H_0^{-1} = 13.5$ Gyr instead of $t_0 = 2/3H_0^{-1} = 8$ Gyr ($\Omega_V = 0$), which is not in conflict with the age of the globular clusters^{19,20}.

7. Bounds on $\tilde{\Omega}_{0V}$

The maximum possible value of $\tilde{\Omega}_{0V}$ is constrained by various bounds produced by the decay of the vacuum and nucleosynthesis^{18,21,22}. The total energy:

$$E = -\ln\left(1 - \tilde{\Omega}_V\right) + V(\phi), \quad (7.1)$$

must decrease between the radiation era and the present epoch giving:

$$1 - \tilde{\Omega}_{0V} > \exp(V_0 - V_R) = \frac{Q_0}{Q_R}, \quad (7.2)$$

where Q_R denotes the value of a quantity at the end of the radiation era.

From nucleosynthesis, we can obtain the inequality:

$$1 - \tilde{\Omega}_{0V} > \frac{1}{(1 + a^2)^{1/2} \xi_{\text{nucleo}}} \approx \frac{1}{\xi_{\text{nucleo}}}, \quad (7.3)$$

where ξ_{nucleo} denotes the speed-up factor: $\xi_{\text{nucleo}} = \tilde{H}_{NS}/\tilde{H}_{\text{standard}}$, where $\tilde{H}_{NS}$ denotes the value of $\tilde{H}$ during nucleosynthesis and $\tilde{H}_{\text{standard}}$ is the value predicted by the standard big-bang model. Because for standard big-bang nucleosynthesis, $\xi_{\text{nucleo}} \sim 1$, this implies that $\tilde{\Omega}_{0V} \approx 0$.

In order to obtain a large enough value of $\tilde{\Omega}_{0V}$ to accomodate the age of the universe, we assume that the coupling of the scalar field ϕ to baryons and anti-baryons is small enough to weaken the bounds on $\tilde{\Omega}_{0V}$ obtained from particle physics and nuclear physics. This will guarantee that the decay of the vacuum and nucleosynthesis do not prevent $\tilde{\Omega}_{0V}$

from reaching a value of order $0.8 - 0.9$, which is needed to obtain a big enough age of the universe.

Why does $\tilde{\Omega}_V$ end up today with the value $\tilde{\Omega}_{0V} \sim 0.8 - 0.9$ needed to give a reasonable age for the universe in the standard model? We can only appeal to the efficiency of the damping mechanism, corresponding to a special potential $V(\phi)$, and to the anthropic principle as answers to this question. It can be argued that given our knowledge of the fundamental constants of nature, the present vacuum density could not be much smaller than it is today to sustain the large-scale structure of the universe^{7,23}.

Acknowledgements

I thank M. Clayton, J. Levin, G. Starkman and D. Yu. Pogosyan for helpful and stimulating discussions. This work was supported by the Natural Sciences and Engineering Research Council of Canada.

References

1. For references, see: P. J. E. Peebles, Principles of Cosmology, Princeton University Press, New Jersey 1993, p.103.
2. M. J. Pierce et al., Nature **371**, 385 (1994).
3. W. L. Freedman et al., Nature, **371**, 757 (1994).
4. A. Guth, Phys. Rev. D **23**, 347 (1981); A. Linde, Rep. Prog. Phys. **47**, 925 (1984).
5. G. Efstathiou et al., Mon. Not. R. Astron. Soc. **247**, 10 (1990); W. Saunders et al., Nature, **349**, 32 (1991).
6. For a review, see: S. M. Carroll, W. H. Press and E. L. Turner, Annu. Rev. Astrophys. **30**, 499 (1992).
7. S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989).

8. P. Jordan, *Nature* **164**, 637 (1949); *Z. Phys.* **157**, 112 (1959).
9. M. Fierz, *Helv. Phys. Acta* **29**, 128 (1956).
10. C. Brans and R. H. Dicke, *Phys. Rev.* **124**, 925 (1961).
11. P. G. Bergmann, *Int. J. Theor. Phys.* **1**, 25 (1968).
12. R. V. Wagoner, *Phys. Rev. D* **1**, 3209 (1970).
13. J. J. Levin and K. Freese, *Phys. Rev. D* **47**, 4282 (1993); *Nucl. Phys. B* **421**, 635 (1994); J. J. Levin, CITA preprints, 1994.
14. V. A. Belinskii et al., *Sov. Phys. JETP* **62**, 195 (1985); V. A. Belinskii and I. M. Khalatnikov, *Sov. Phys. JETP* **66**, 441 (1987).
15. D. La and P. J. Steinhardt, *Phys. Rev. Lett.* **62**, 376 (1989).
16. J. D. Barrow and Kei-ichi Maeda, *Nucl. Phys. B* **341**, 294 (1990).
17. S. Kalara, N. Kaloper and K. A. Olive, *Nucl. Phys. B* **341**, 252 (1990).
18. T. Damour and K. Nordtvedt, *Phys. Rev. D* **48**, 3436 (1993).
19. P. Demarque, C. P. Deliyannis and A. Sarajedini, *Observational Tests of Cosmological Inflation*, ed. T. Shanks et al., 111. Dordrecht: Kluwer (1991).
20. B. Chaboyer, CITA preprint, 1994.
21. M. Öser and M. O. Taha, *Phys. Lett.* **171**, 363 (1986); K. Freese et al., *Nucl. Phys. B* **287**, 797 (1987).
22. B. Ratra and P. J. E. Peebles, *Phys. Rev. D* **37**, 3406 (1988).
23. G. Starkman, in preparation.